



Anthelmintic effects of sa infoin against different catt le nematodes may be linked to concentration of condensed tannins in different gut compartments

Desrues, Olivier; Pena-Espinoza, Miguel Angel; Hansen, Tina V.A. ; Mueller-Harvey, Irene; Enemark, Heidi; Thamsborg, Stig M.

Published in:

25th International Conference of the World Association for the Advancement of Veterinary Parasitology

Publication date:

2015

Document Version

Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):

Desrues, O., Pena-Espinoza, M. A., Hansen, T. V. A., Mueller-Harvey, I., Enemark, H., & Thamsborg, S. M. (2015). Anthelmintic effects of sa infoin against different catt le nematodes may be linked to concentration of condensed tannins in different gut compartments. In *25th International Conference of the World Association for the Advancement of Veterinary Parasitology: Abstract Book* (pp. 58-58).

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Monday 17 August 2015

Concurrent Session 1.3

C: Novel approaches to chemical and non-chemical control (endo and ecto-parasites)

Ref: O018/0628

Anthelmintic effects of sainfoin against different cattle nematodes may be linked to concentration of condensed tannins in different gut compartments

Olivier Desrues¹, Miguel Peña-Espinoza², Tina V. Hansen¹, Irene Mueller-Harvey⁴, Heidi L. Enemark^{2,3}, Stig M. Thamsborg¹

¹Department of Veterinary Disease Biology, University of Copenhagen, Frederiksberg C, Denmark, ²National Veterinary Institute, technical University of Denmark, Frederiksberg C, Denmark, ³Norwegian Veterinary Institute, Oslo, Norway, ⁴School of Agriculture, Policy and Development, University of Reading, Reading, UK

Objectives: We investigated the potency of a forage legume (sainfoin, *Onobrychis viciifolia*) against gastrointestinal nematodes in relation to concentration of condensed tannins (CT) along the digestive tract in cattle.

Methods: Fifteen 2-4 month-old Jersey calves were divided into two groups and fed either sainfoin pellets (SF, n=9) or starter feed pellets and ryegrass hay (CO, n=6). Diets were balanced for crude protein and net energy during the entire study. Then, calves were infected with 10,000 *Ostertagia ostertagi* and 66,000 *Cooperia oncophora* third-stage larvae (L3). The number of eggs per g of dried faeces (FECDM) was followed until 42 days post infection when the calves were euthanized for worm recovery and sampling of digesta for analysis of CT. The HCl-butanol-acetone and *in situ* thiolysis methods were used to assess the CT content and structural composition.

Results: The main finding was a 50 % reduction in adult worm counts for *O. ostertagi* in SF as compared to CO ($p < 0.05$). In contrast no difference was observed between groups for *C. oncophora*. Overall, FECDM was 40% lower in SF compared to CO ($p > 0.05$). The CT content was 6.5% of dry matter (DM) in sainfoin pellets and gradually increased along the digestive tract (5.9, 6.6 and 8.1% DM in the abomasum, small intestine and faeces, respectively). However, *in situ* thiolysis indicated more CT in the abomasum and less CT in other locations of the digestive tract. The CT contents for pellets, abomasum, small intestine and faeces were 1.9, 2.2, 0.01 and 0.5% DM, respectively. This suggests that CT become less accessible and reactive to the thiolysis reagent as the feed passes along the digestive tract.

Discussion: Feeding sainfoin significantly reduced the worm burden of *O. ostertagi* in the abomasum where CT are thought to be released from complexes with macromolecules due to acidic conditions. The lack of effect against *C. oncophora* in the small intestine could be related to an apparent unavailability of CT in this gut compartment.